

Molecular reactions of PAH fragments in astronomical environments irradiated by soft X-rays

Sergio A. Cifuentes-Vásquez^{1*}, Heidi M. Qutián-Lara² & Mario A. Higuera-Garzón³

¹Universidad Nacional de Colombia, Facultad de Ciencias, Departamento de Química

²Max-Planck-Institute für Extraterrestrische Physik, Center for Astrochemical Studies

³Universidad Nacional de Colombia, Facultad de Ciencias, Observatorio Astronómico Nacional

*scifuentesv@unal.edu.co

VIII Congreso Colombiano de Astronomía y Astrofísica, CoCoA2024

Table of contents

01

Introduction

PAHs and prebiotic
molecules in space

02

Data acquisition

MS-TOF
Sigma-Aldrich reagents

03

Results

What has been achieved?

04

Future perspectives

What is yet to come?

05

Acknowledgements

06

Bibliography

01

Introduction


PAHs and prebiotic molecules in space

Introduction

Prebiotic molecules, predominantly organic, are considered pivotal in the origin of life on Earth and potentially in other planetary environments.


Relevant molecules:

- Aromatic hydrocarbons
- Ammonia derivatives
- Carbon compounds




Introduction

In circumstellar regions, benzenoid molecules undergo significant modifications under the influence of ultraviolet (UV) photons and X-rays. Processes such as ionization, dissociation, and photo-fragmentation result in the formation of its ionic species and free radicals, potentially leading to the creation of new organic molecules and giving valuable insights into mechanisms.




Objective


Investigate the impact of soft X-rays on the physicochemical properties of aromatic hydrocarbons, their potential involvement in the genesis of prebiotic molecules, and their potentially interesting applications in storage devices.



benzene



benzene- d_6




2-methylantracene

02

Data acquisition

MS-TOF

Sigma-Aldrich reagents





- Samples were purchased from Sigma-Aldrich
- Radiation source: Spherical Grating Monochromator (SGM)
- Ultrahigh vacuum chamber: $\approx 10^{-8}$ mbar
- Drift tube: 297 mm

03



Results

What has been achieved?



Results for benzene





Results for benzene



Results for benzene





Results for benzene-d **6**





Results for benzene-d ₆

6





Results for benzene-d ₆

6



Preliminary results for 2-methylantracene





04

Future perspectives

What is yet to come?

Future perspectives

The resulting PAH fragments product of the soft X-ray fragmentation and photo-dissociation may interfere in the formation of diverse COMs. They may also give valuable insights about the genesis of prebiotic molecules as well as other applications relevant to modern investigations about new materials or new appliances subjected to harsh and hostile conditions such as astronomical environments.



05

Acknowledgments

Acknowledgements

The Universidad Nacional de Colombia supported this work via project QUIPU 202010042199, and Min-Ciencias through Conv. 937.



06

Bibliography

Bibliography

- [1] Boechat-Roberty, H. M.; Neves, R.; Pilling, S.; Lago, A. F.; de Souza, G. G. Dissociation of the Benzene Molecule by Ultraviolet and Soft X-Rays in Circumstellar Environment. *Monthly Notices of the Royal Astronomical Society* 2009, 394 (2), 810–817.
- [2] Higuera-G., M.-A.; Ramos P., A. F. PAHs as tracers of local AGN-starburst connection. *Revista Mexicana de Astronomía y Astrofísica*. 2013, 49, 301–309.
- [3] Huo, Y.; Cangahuala, M. K.; Zamudio-Bayer, V.; Goulart, M.; Kubin, M.; Timm, M.; Lau, J. T.; Issendorff, B. von; Hoekstra, R.; Faraji, S.; Schlathölter, T. A Comparative Laboratory Study of Soft X-Ray-Induced Ionization and Fragmentation of Five Small PAH Cations. *The European Physical Journal D* 2023, 77 (10).
- [4] Quitián-Lara, H. M.; Fantuzzi, F.; Oliveira, R. R.; Nascimento, M. A.; Wolff, W.; Boechat-Roberty, H. M. Dissociative Single and Double Photoionization of Biphenyl (C₁₂H₁₀) by Soft x-Rays in Planetary Nebulae. *Monthly Notices of the Royal Astronomical Society* 2020, 499 (4), 6066–6083.
- [5] Ramos Padilla, A. F.; Ashby, M. L. N.; Smith, H. A.; Martínez-Galarza, J. R.; Beverage, A. G.; Dietrich, J.; Higuera-G., M.-A.; Weiner, A. S. The AGN contribution to the UV-FIR luminosities of interacting galaxies and its role in identifying the main sequence. *Monthly Notices of the Royal Astronomical Society* 2020, 499 (4), 4325–4369.

Thank you for your attention!

CREDITS: This presentation template was created by [Slidesgo](#), and includes icons by [Flaticon](#), and infographics & images by [Freepik](#)